

**D.K.M.COLLEGE FOR WOMEN (AUTONOMOUS), VELLORE-1.**

**COMPLEX ANALYSIS**

**Unit-I      SECTION-A      2 Marks**

1. Define Triangle inequality.
2. Prove that  $|z - 1| = |z + i|$  represents the line through the origin  
Origin whose slope is -1
3. The limit of a function is unique.
4. Define entire function.
5. Define harmonic function.
6. Prove that  $u = y^3 - 3x^2y$  is harmonic
7. If  $f'(z) = 0$  in a domain D. then  $f(z)$  must be a constant throughout D
8. Define analytic function.

**SECTION-B      5 Marks**

1. Suppose that  $f(z) = u(x, y) + iv(x, y)$ ,  $z_0 = x_0 + iy_0$ ,  $w_0 = u_0 + iv_0$  then  
 $\lim_{z \rightarrow z_0} f(z) = w_0$  if  $\lim_{(x, y) \rightarrow (x_0, y_0)} u(x, y) = u_0$ ,  $\lim_{(x, y) \rightarrow (x_0, y_0)} v(x, y) = v_0$
2. Explain stereographic projection.
3. If the derivatives are  $f$  and  $F$  exists at a points  $Z$  then
  - i.  $(f \pm F)'(z) = f'(z) \pm F'(z)$
  - ii.  $(f \cdot F)'(z) = f(z)F'(z) + f'(z)F(z)$

$$\text{iii. } \left(\frac{f}{F}\right)'(z) = \frac{F(z)f'(z) - F'(z)f(z)}{(F(z))^2}$$

Provided  $F(z) \neq 0$

4. Necessary condition for a function  $f$  to be differentiable at  $z_0$ . Suppose that  $f(z) = u(x, y) + iv(x, y)$  and  $f'(z)$  exist at  $z_0 = x_0 + iy_0$  then the first order partial derivatives of  $u$  and  $v$  must exist at  $(x_0, y_0)$  and they satisfy Cauchy-Riemann equations.  $u_x = v_y$ ,  $u_y = -v_x$  and  $f'(z)$  can be written as  $f'(z_0) = u_x + iv_x$ . Where this partial derivatives are to be evaluated at  $(x_0, y_0)$ .

$$5. \text{ For any function of } \phi \frac{\partial^2 \phi}{\partial x^2} + \frac{\partial^2 \phi}{\partial y^2} = 4 \frac{\partial^2 \phi}{\partial z \partial \bar{z}}$$

6. An analytic function with constant modulus is constant.

7. If  $f(z) = u + iv$  is an analytic form of  $z$  and  $u - v = \frac{\cos x + \sin x - e^{-y}}{2 \cos x - e^y - e^{-y}}$  find  $f(z)$  subject to the condition  $f\left(\frac{\pi}{2}\right) = 0$

### SECTION-C

**10 MARKS**

1. If  $f$  is differentiable at  $z_0$ , then  $f$  is continuous at  $z_0$
2. Sufficient condition for a function  $f(z)$  to be differentiable at  $z_0$ .
3. Let the function  $f(z) = u(r, \theta) + iv(r, \theta)$ . We define throughout some  $\varepsilon$ -neighbourhood of a non-zero point  $z_0 = r_0 \exp(i\theta_0)$
4. If  $f(z)$  is analytic function of  $z$   
Prove that  $\left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2}\right) = 4 \left(\frac{\partial^2}{\partial z \partial \bar{z}}\right)$   
Prove that  $\left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2}\right) |f(z)|^2 = 4 |f'(z)|^2$
5. Necessary condition for a function  $f$  to be differentiable at  $z_0$ .
6. If  $\lim_{z \rightarrow z_0} f(z) = w_0, \lim_{z \rightarrow z_0} f'(z) = W_0$

Then

$$\text{i. } \lim_{z \rightarrow z_0} f(z) \pm F(z) = w_0 \pm W_0$$

- ii.  $\lim_{z \rightarrow z_0} f(z) \cdot F(z) = w_0 W_0$
- iii.  $\lim_{z \rightarrow z_0} f(z)/f(z) = \frac{w_0}{W_0}$ , where  $w_0 \neq 0$

### UNIT -II

### SECTION-A

2 MARKS

1.  $\omega = b + Az$  where A & B are complex constant.
2. Find the image strip  $0 < x < 1$  under the transformation  $w = iz$
3. Define isogonal and conformal transformation
4. Define critical point .with example
5. Define  $w = z^2$

### SECTION-B

5 MARKS

1. Every bilinear transformation preserves the cross ratio
2. Find the bilinear transformation with maps  $z_1 = -1, z_2 = 0, z_3 = 1$  to  $w_1 = -1, w_2 = 1, w_3 = i$
3. Every bilinear transformation which has only one fixed point can be put in the form  $\frac{1}{w - \alpha} = \frac{1}{z - \alpha} + \lambda$
4. Find the fixed point and bring it to the normal form of following bi-linear transformation.  $\omega = \frac{(z+i)z-2}{z+i}$
5. Find the bi-linear transformation which transformation the half plane  $\text{Re}(z) \geq 0$  into the circle  $|\omega| \leq 1$
6. Find the mobius transformation which transformation the circle  $|z| = 1$  and  $|\omega| = 1$  and makes the points  $z = 1, -1$  correspondence to  $\omega = 1, -1$

### SECTION-C

10 MARKS

1. Discuss the transformation  $\omega = \frac{1}{z}$
2. Necessary condition for conformality.
3. Sufficient condition for conformality.
4. Discuss the transformation  $\omega = e^z$
5. Discuss the transformation  $\omega = z^2$
6. Find the bilinear transformation which transforms the circle

$|z| \leq \rho$  to the circle of  $|\omega| \leq \rho$

**UNIT III**

**SECTION-A**

**2 MARKS**

1. Define arc and simple arc.
2. Define simple closed curve and Jordan curve .
3. Find the value of integral  $I = \int_c \bar{z} dz$  where  $c$  is the right hand half  $z = 2e^{i\theta}$   $\left(-\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}\right)$  of the circle  $|z| = 2$  from  $z = -2i$  to  $z = 2i$
4. Define multiple connected domain.
5. Define Cauchy-goursat for multiple connected domain.

**SECTION-B**

**5 MARKS**

1.  $C_1$  denote the contour OAB find  $\int_c f(z) dz$  where  $f(z) = y - x - i3x^2$ ,  $O(0,0)$ ,  $A(i)$ ,  $B(1+i)$
2.  $I = \int_c \frac{1}{z^2} dz$  where  $c$  is the semicircle  $z = 3e^{i\theta}$  where  $0 \leq \theta \leq \pi$  from the point  $z = 3$  to  $z = -3$

**SECTION-C**

**10 MARKS**

1. State and prove cauchy's theorem.
2. If a function  $f$  is analytic at a point then its derivatives of all orders at that point and they are analytic.
3. State and prove maximum modulus theorem.
4. Suppose that  $f(z)$  is analytic throughout a neighbourhood  $|z - z_0| < \epsilon$  of a point  $z_0$ . If  $|f(z)| \leq |f(z_0)|$  for each point  $z$  in that neighbourhood then  $f(z)$  has a constant value  $f(z_0)$  throughout the neighbourhood.
5. State and prove maximum Cauchy integral formula.

**UNIT IV****SECTION-A****2 MARKS**

1. Define convergent and divergent series.
2. Define maclaurin series.
3. Define pole.
4. Define essential singular point and removable singular point
5. Define principal part of f at  $z_0$ .

**SECTION-B****5 MARKS**

1. Suppose that  $z_n = x_n + iy_n$ ,  $n=1,2,\dots, z=x+iy$  then  
 $\lim_{n \rightarrow \infty} z_n = z \Leftrightarrow \lim_{n \rightarrow \infty} x_n = x ; \lim_{n \rightarrow \infty} y_n = y$ .
2. Suppose that  $z_n = x_n + iy_n$ ,  $n=1,2,\dots, z=x+iy$  then  
 $\sum_{n=1}^{\infty} x_n = X$  and  $\sum_{n=1}^{\infty} y_n = y$
3.  $F(z) = \frac{1}{z(z^2 - 3z + 2)}$  for the region
  - i.  $0 < |z| < 1$
  - ii.  $1 < |z| < \infty$
  - iii.  $|z| > 1$

**SECTION-C****10 MARKS**

1. State and prove Taylor's series.
2. State and prove Laurent's series.

**UNIT V****SECTION-A****2 MARKS**

1. Find the poles and residues  $f(z) = \frac{1}{z+z^2}$
2. Find the poles and residues  $f(z) = \frac{1}{z^2+a^2}$

**SECTION-B****5 MARKS**

1. State and prove Cauchy residue theorem.

2. An isolated singular point  $z_0$  of a function  $f$  is a pole of order  $m$  iff  $f(z) = \frac{\phi(z)}{(z-z_0)^m}$  where  $\phi(z)$  is analytic and non-zero at  $z_0$  and if  $m=1$   $\text{res}_{z=z_0} f(z) = \phi(z_0)$  and if  $m \geq 2$   $\text{res}_{z=z_0} f(z) = \frac{\phi^{(m-1)}(z_0)}{(m-1)!}$
3.  $\int_c \frac{z+1}{z^2-2z} dz$  where  $c$  is the circle  $|z| = 3$
4. find the pole and residue  $f(z) = \left(\frac{z}{2z+1}\right)^3$
5. find the pole and residue  $f(z) = \frac{1}{z^2+a^2}$

### SECTION-C

10 MARKS

1. Show that  $\int_0^\infty \frac{x^2 dx}{(x^2+1)(x^2+4)} = \frac{\pi}{6}$
2. Show that  $\int_{-\infty}^\infty \frac{\cos 3x dx}{(x^2+1)^2} = \frac{2\pi}{e^3}$